Sanitized Copy Approved for Release 2011/08/31: CIA-RDP80-00809A000600330562-1

DEC. 1948

SECRET

CLASSIFICATION SECRET

CENTRAL INTELLIGENCE AGENCY

REPORT

50X1-HUM

INFORMATION FROM FOREIGN DOCUMENTS OR RADIO BROADCASTS

CD NO. ST-74

COUNTRY

USSR

Scientific - Medical hemorrhagic fever

DATE OF

INFORMATION 1948

SUBJECT

DATE DIST. # Aug 1950 .

HOW

PUBLISHED Bimonthly periodical

PUBLISHED MOSCOW

NO. OF PAGES

DATE

PUBLISHED Apr 1948

SUPPLEMENT TO REPORT NO.

LANGUAGE Russian

THIS DOCUMENT CONTAINS INFORMATION AFFECTING THE MATIONAL DEFEN OF THE UNITED STATES WITHIN THE MEANING OF REPIONALE ACT I U. S. C., 31 AND 32, AS ANENDED. ITS TRANSMISSION OF ATHE REVELATION OF TIS CONTENTS IN ANY MARKER TO AN UNAUTHORIZED PERSON IS PR MISITED BY LAW. REPRODUCTION OF THIS FORM IS PROMISITED.

THIS IS UNEVALUATED INFORMATION

SOURCE

Vestnik Akademii Meditsinskikh Nauk SSR, No 2, 1948.

EXPEDITIONS FOR THE STUDY OF OMSK EPIDEMIC HEMORRHAGIC FEVER

∠A Digest7

During the past few years, the Institute of Neurology has activated an expedition for the study of Omsk hemorrhagic fever (OL), a new infectious disease, of unknown etiology, attacking man in the Central Asiatic regions of the USSR.

As early as 1944, members of the Academy of Medical Sciences USSR and the then independent Primorskaya Army in Crimea were conducting studies on Crimean hemorrhagic fever in an attempt to define the immunological, etiological, epidemiological, and serological characteristics of this infection. Results of these studies were published in 1945, in Crimean Hemmorrhagic Fever, and in 1947, in the series Medical News and the volume Virus Diseases.

In 1946, an expedition was sent into the Omsk region to make a study of an epidemic disease whose characteristics were very similar to those of Crimean hemorrhagic fever. The absence of the tick Hyaloma marginatum, carrier of Crimean hemorrhagic fever, in the fuana of Omsk Oblast made studies difficult. The studies were carried out under the supervision of Prof R. M. Akhremovich and several members of the Omsk Medical Institute including G. A. Sizemova, Yu. V. Veselova, V. P. Konstantinov, I. S. Novitskiy, A. A. Fedyushin, entomologist G. I. Netskiy, and others. No definite results were obtained by this expedition, with the result that in 1947 the Omsk Oblast Public Health Department of the Ministry of Public Health USSR and the Academy of Medical Sciences organized a special expedition to study OL.

This expedition was placed under the supervision of Prof M. P. Chumakov, Laureate of the Stalin Prize, and included the following members: Prof A. F. Bilibin, Yu. S. Kleyn, neuropathologist Dr. N. V. Sorokina, virusologist and Scientific Associate A. P. Belyayeva, N. S. Slavina, A. V. Gagarina, and Laboratory technicians O. A. Arapova and K. D. Vinogradova. In addition, workers were recruited from clinics of medical institutes, the Oblast Institute of Microbiology, malarial stations, and the Agricultural Institute. The expedition consisted of a total of 50 members. The virusological group, which was also under the supervision of Prof M. P. Chumakov, was divided into two clinical groups headed by Prof A. Bilibin and Prof R. M. Akhremovich. Three other groups contained within the expedition included the parasitological groups,

CLASSIFICATION	SECRET	 	
STATE X NAVY X NSRB	DISTRIBUTION	 +	
ARMY X AIR X FBI		,	

CL.	ΛD	CT
SE	UN	E.

SECRET

50X1-HUM

consisting of Prof A. V. Fedyushin and entomologist G. I. Netskiy; the epidemiological group headed by Docent M. V. Los; and the pathologoanatomical group under Prof I. S. Novitskiy.

In addition to research conducted by the expedition, M. P. Chumakov and Dr A. P. Belyayeva conducted laboratory experiments in their laboratory at the Institute of Neurology, Academy of Medical Sciences USSR, to determine the effects of filtrates obtained from human poliomyelitic victims on monkeys. Experiments on immunity were conducted on monkeys at the Sukhumi Biological Staperiments of the Academy of Medical Sciences USSR by Dr S. Skokorshumova of the Institute of Virusology, M. K. Voroshilova, and L. I. Mart'yanova.

In 1945 and 1946, associates of the Omsk Institute of Microbiology and Tularemia Station, including A. A. Gavrilovskaya, Ye. D. Kushnir, and others, conducted detailed studies to determine the microbe agent in the etiology of OL. The steppe tick Dermacentor pictus was found to contain OL virus, and thus might be the source of infection in humans. It was possible to determine a method for serodiagnosis in isolated cases of OL.

As a result of data collected by the Chumakov Expedition it was possible to make the following statements:

- 1. The causative agent of OL was found to be a filterable virus, present in the blood of patients and pathogenic to many laboratory animals. The etiological role of this virus, which is passive on animals, has been defined, in view of the regularity of its detection in patients, the presence of specific antibodies against this virus in RSK (complement fixation reaction) and RN antibodies against this virus in RSK (complement fixation reaction) and RN in view of the identical clinical aspects of naturally developed infections of OL and accidental laboratory infections occasioned after working with this virus.
- Valuable materials were collected on the clinical, laboratory, pathomorphological, and epidemiological characteristics of OL.
- 3. Steppe ticks Dermacentor pictus, collected in the OL endemic regions contained virus and hence may be carriers.
- 4. The development of an immunity to OL in those vaccinated with killed virus established the possibility of practical use of vaccination in combating this fever.
- 5. In tests on monkeys, mice and other animals, it was shown that there was an absence of crossed immunological reactions between OL and Crimean hemorrhagic fever as well as an absence of etiological independence of each of these infections, which in clinical-epidemiological relationship appeared only similar but not analogous to each other. On the basis of experiments with crossed immunity on guinea pigs and as a result of regular morphological studies it was possible to differentiate the OL virus from the filtered, rickettsial type of Q fever, or the so-called atypical virus pneumonia.
- 6. The possibility of practical serodiagnosis of individual cases of OL and those suspected of OL illnesses was established by means of reaction of uniting the complement and reaction due to neutralling the virus with specific serums.

~ E N D ~

- 2 -

SECRET SECRE